

Ordered valence bond states in symmetric two-dimensional spin-orbital systems

Guang-Ming Zhang^{1,2} and Shun-Qing Shen¹

¹*Department of Physics, the University of Hong Kong, Pokfulam Road, Hong Kong, China.*

²*Center for Advanced Study, Tsinghua University, Beijing, 100084, China*

(February 1, 2008)

We consider a superexchange Hamiltonian, $H = -\sum_{\langle i,j \rangle} (2\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{2})(2\mathbf{T}_i \cdot \mathbf{T}_j - \frac{1}{2})$, which describes systems with orbital degeneracy and strong electron-phonon coupling in the limit of large on-site repulsion. In an $SU(4)$ Schwinger boson representation, a reduced spin-orbital interaction is derived *exactly*, and a mean field theory has been developed by introducing a symmetric valence bond pairing order parameter. In one dimension, a spin-orbital liquid state with a finite gap is obtained. On a two-dimensional square lattice a novel type of spin-orbital ferromagnetically ordered state appears, while spin and orbital are antiferromagnetic. Moreover, an important relation has been found, relating the spin and orbital correlation functions to the combined spin-orbital ones.

PACS numbers: 75.10.Jm, 11.10.Hi, 11.25.Hf, 75.40.Mg

There have been much interests in the properties of Mott insulators with orbital degeneracy [1–4], where the electron configuration has an orbital degeneracy in addition to the spin degeneracy. Due to the interplay between the spin and orbital degrees of freedom, a rich variety of spin and orbital ordering effects have been displayed [5–9], and a new combined spin-orbital degrees of freedom may introduce some new physics to the transitional metal oxides [10–15]. Recently, in order to describe the low-energy physics of an insulating crystal with one-electron per site with double orbital degeneracy in the limit of the large on-site Hubbard repulsion, a superexchange Hamiltonian was proposed [16,17]

$$H = - \sum_{\langle i,j \rangle} (2\mathbf{S}_i \cdot \mathbf{S}_j - \frac{1}{2})(2\mathbf{T}_i \cdot \mathbf{T}_j - \frac{1}{2}), \quad (1)$$

where two isotropic Heisenberg antiferromagnets (AFM) coupled by a quartic term on equivalent bonds, and $\mathbf{S}_i$ and $\mathbf{T}_i$ denote the spin-1/2 and orbital-1/2 operators at a lattice site i , respectively. The physical condition to derive the above effective model Hamiltonian is that among the possible two-particle states obtained upon virtual hopping, the inter-orbital singlet is the lowest energy state due to the dynamical Jahn-Teller effect. The condition might be realized in the large body of new molecular compounds based on C_{60} [18] or layered fullerenes and some two-dimensional copolymers, such as vinylidene fluoridetrifluoroethylene [19]. Based on numerical calculations, Santoro *et al.* argued that the ground state of the model shows a spin-Peierls-like dimerization in one dimension (1D) [16], while on a two dimensional (2D) square lattice no evidence of order of any kinds had been found that the ground state is concluded to be a spin liquid of resonating valence bonds (VB) [17]. In order to put these results on a firm ground, it is thus of great interest to develop new approaches to this model.

In this Letter, we use an $SU(4)$ Schwinger boson representation [20] to denote both the spin-1/2 and orbital-1/2 operators at the same time, and the model Hamiltonian is simplified to a reduced interaction, describing an

symmetric pairing attraction among the nearest neighbor hard-core bosons. By introducing an symmetric VB order parameter, a mean field theory is developed. In 1D, a spin-orbital liquid state with a finite gap in the excitation spectra is obtained, corresponding to a quantum *disordered* VB state. On a 2D square lattice, due to the presence of the Bose-Einstein condensations we found a novel spin-orbital ferromagnetically (FM) ordered state, corresponding to a short-ranged VB crystal state, while both the spin and orbital degrees of freedom form an AFM ordering.

The model Hamiltonian can be rewritten as follows

$$H = 2 \sum_{\langle i,j \rangle} (\mathbf{S}_i \cdot \mathbf{S}_j + \mathbf{T}_i \cdot \mathbf{T}_j) - \sum_{\langle i,j \rangle} (2\mathbf{S}_i \cdot \mathbf{S}_j + \frac{1}{2})(2\mathbf{T}_i \cdot \mathbf{T}_j + \frac{1}{2}), \quad (2)$$

where the direct quadratic couplings among spins and orbitals are antiferromagnetic, respectively, while the quartic coupling between spins and orbitals is ferromagnetic. First of all, the model Hamiltonian has an $SU(2) \otimes SU(2)$ symmetry, representing rotational invariance in both spin and orbital space, and also interchange symmetry between the spins and orbitals. Moreover, one also notices that the total spin, orbital, and combined *staggered* spin-orbital operators

$$S^\alpha = \sum_j S_j^\alpha, \quad T^\alpha = \sum_j T_j^\alpha, \quad L^{\alpha\beta} = \sum_j e^{i\mathbf{Q} \cdot \mathbf{R}_j} 2S_j^\alpha T_j^\beta,$$

where $\mathbf{Q}$ is the AF reciprocal vector, generate an $SU(4)$ Lie algebra, and also commutes with the model Hamiltonian. Therefore, Eq.(1) possesses the $SU(4)$ symmetry [16]. For a state with the $SU(4)$ symmetry, implying that the state is invariant under the rotation in the $SU(4)$ space, we have an identity for the static correlation functions

$$\langle S_i^\alpha S_j^\alpha \rangle = \langle T_i^\beta T_j^\beta \rangle = e^{i\mathbf{Q} \cdot (\mathbf{R}_i - \mathbf{R}_j)} \langle 4S_i^\alpha T_i^\beta S_j^\alpha T_j^\beta \rangle, \quad (3)$$

where $\langle \dots \rangle$ represents the expectation value on the SU(4) symmetric state. The underlying physics for the identities is very clear: if the spin and orbital degrees of freedom are antiferromagnetically correlated, the combined spin-orbital correlation functions will be ferromagnetic. This property originates from the additional phase factor in the generators $L^{\alpha\beta}$. It should be mentioned that another spin-orbital model Hamiltonian $H' = \sum_{\langle i,j \rangle} (2\mathbf{S}_i \cdot \mathbf{S}_j + \frac{1}{2})(2\mathbf{T}_i \cdot \mathbf{T}_j + \frac{1}{2})$, the natural generalization of the SU(2) Heisenberg spin model and *exactly* soluble in 1D, has also been used to understand some orbital properties of transition metal oxides [10–15]. However, although these two models have the same symmetry, the fifteen generators of the SU(4) symmetry group are not the same, and the identities of the correlation functions induced from this symmetry are also different. In this sense, the physics involved in these two models are independent, leading to different spin-orbital properties of transitional metal oxides.

As we know, the Schwinger boson mean field approach has been a successful theory in describing the low-energy excitations of the conventional FM and AFM Heisenberg spin superexchange models [20,21]. In particular, starting from short-ranged VB order parameters, it can produce either quantum disordered or quantum long-range ordered magnetic states. In the present model when two sets of SU(2) Schwinger bosons are introduced to denote the spin-1/2 and orbital-1/2 operators separately, we have

$$2\mathbf{S}_i \cdot \mathbf{S}_j + \frac{1}{2} = \sum_{\alpha,\beta} a_{i,\alpha}^\dagger a_{i,\beta} a_{j,\beta}^\dagger a_{j,\alpha}$$

$$2\mathbf{T}_i \cdot \mathbf{T}_j + \frac{1}{2} = \sum_{m,n} d_{i,m}^\dagger d_{i,n} d_{j,n}^\dagger d_{j,m}$$

with local constraints $\sum_{\alpha} a_{i,\alpha}^\dagger a_{i,\alpha} = 1$ and $\sum_m d_{i,m}^\dagger d_{i,m} = 1$, where the indices α, β and m, n are taken values 1 and 2 corresponding to eigenstates of $S^z = \pm 1/2$ and $T^z = \pm 1/2$, respectively. Both $a_{i,\alpha}$ and $d_{i,m}$ operators satisfy the boson commutation relations. To treat the quartic spin-orbital interaction term on an equal footing as the same as the spin-spin and orbital-orbital superexchange interactions, the fact has been used that in the model Hamiltonian the Hilbert space on each lattice site consists of four basic states in terms of $|S^z; T^z\rangle$:

$$\left| +\frac{1}{2}; +\frac{1}{2} \right\rangle = |1\rangle, \quad \left| -\frac{1}{2}; +\frac{1}{2} \right\rangle = |2\rangle,$$

$$\left| +\frac{1}{2}; -\frac{1}{2} \right\rangle = |3\rangle, \quad \left| -\frac{1}{2}; -\frac{1}{2} \right\rangle = |4\rangle.$$

These four states form a set of local basis to represent the SU(4) symmetry group. The conventional SU(4) generators $S_\mu^\nu(i)$ act on a basic state $|\eta\rangle_i$ according to the equation $S_\mu^\nu(i)|\eta\rangle_i = \delta_{\nu,\eta}|\mu\rangle_i$ with a local constraint $\sum_\mu S_\mu^\mu(i) = 1$, where the indices μ and ν are taken values from 1 to 4 corresponding to the four eigenstates of

$|\pm\frac{1}{2}; \pm\frac{1}{2}\rangle$. The SU(4) Lie algebra is defined by the commutation relation

$$[S_m^n(i), S_k^l(i)] = \delta_{n,k} S_m^l(i) - \delta_{m,l} S_k^n(i).$$

In terms of four-component hard-core bosons, we have $S_\mu^\nu(i) = b_{i,\mu}^\dagger b_{i,\nu}$ with the local constraint $\sum_\mu b_{i,\mu}^\dagger b_{i,\mu} = 1$. Therefore, the quartic spin-orbital exchange interaction can be written as

$$(2\mathbf{S}_i \cdot \mathbf{S}_j + \frac{1}{2})(2\mathbf{T}_i \cdot \mathbf{T}_j + \frac{1}{2}) = \sum_{\mu,\nu} b_{i,\mu}^\dagger b_{i,\nu} b_{j,\nu}^\dagger b_{j,\mu}, \quad (4)$$

By a projection procedure, both the spin-spin and orbital-orbital quadratic superexchange interactions can also be expressed in terms of the four-component hard-core bosons. Fortunately, most of the resulting terms are found to be *exactly* canceled with the quartic spin-orbital exchange interaction, and the resulting Hamiltonian is reduced to a simple and compact form

$$H = - \sum_{\langle i,j \rangle} \left[(b_{i,1}^\dagger b_{j,4}^\dagger + b_{i,4}^\dagger b_{j,1}^\dagger) - (b_{i,2}^\dagger b_{j,3}^\dagger + b_{i,3}^\dagger b_{j,2}^\dagger) \right] \\ \times [(b_{j,4} b_{i,1} + b_{j,1} b_{i,4}) - (b_{j,3} b_{i,2} + b_{j,2} b_{i,3})], \quad (5)$$

Actually, this reduced model Hamiltonian has explicitly displayed that the main physics of the symmetrically coupled spin-orbital interaction is to *induce a unique attractive pairing instability among the nearest neighbor hard-core bosons*. We would also emphasize here that the quartic spin-orbital exchange interaction plays the *same* role as that of the spin-spin and orbital-orbital superexchange terms, as both quadratic and quartic exchange interactions have similar forms and are treated on the equal footing.

In the reduced spin-orbital model Hamiltonian, the local attractive interaction shares closely resemblance with the effective model Hamiltonian in the conventional BCS superconductivity theory. Such a feature allows us to introduce a unique short-ranged VB pairing order parameter

$$\Delta = -\langle (b_{j,4} b_{i,1} + b_{j,1} b_{i,4}) - (b_{j,3} b_{i,2} + b_{j,2} b_{i,3}) \rangle,$$

which can assume to be real. A mean field theory is followed to develop naturally, leading to the mean field Hamiltonian

$$H_{mf} = \lambda \sum_{i,\mu} b_{i,\mu}^\dagger b_{i,\mu} - \lambda N + Z \Delta^2 N \\ + \Delta \sum_{\langle i,j \rangle} \left[(b_{i,1}^\dagger b_{j,4}^\dagger + b_{i,4}^\dagger b_{j,1}^\dagger) - (b_{i,2}^\dagger b_{j,3}^\dagger + b_{i,3}^\dagger b_{j,2}^\dagger) + h.c. \right],$$

where the local constraint has been imposed on average through a Lagrangian multiplier, N is the total number of lattice sites, and Z is the number of the nearest neighbors. In the momentum space, H_{mf} becomes

$$H_{mf} = \lambda \sum_{\mathbf{k}, \mu} b_{\mathbf{k}, \mu}^\dagger b_{\mathbf{k}, \mu} - \lambda N + Z \Delta^2 N + 2Z \Delta \sum_{\mathbf{k}} \gamma_{\mathbf{k}} \left[(b_{\mathbf{k}, 1}^\dagger b_{-\mathbf{k}, 4}^\dagger - b_{\mathbf{k}, 2}^\dagger b_{-\mathbf{k}, 3}^\dagger) + h.c. \right], \quad (6)$$

where $\gamma_{\mathbf{k}} = \frac{1}{Z} \sum_{\delta} e^{i\mathbf{k} \cdot \delta}$ and the vector δ points to the nearest neighbor sites. When a Nambu spinor is defined by $\Psi^\dagger(\mathbf{k}) = (b_{\mathbf{k}, 1}^\dagger, b_{\mathbf{k}, 2}^\dagger, b_{\mathbf{k}, 3}^\dagger, b_{\mathbf{k}, 4}^\dagger; b_{-\mathbf{k}, 4}, b_{-\mathbf{k}, 3}, b_{-\mathbf{k}, 2}, b_{-\mathbf{k}, 1})$, the mean field Hamiltonian is expressed in a compact matrix form

$$H_{mf} = \frac{1}{2} \sum_{\mathbf{k}} \Psi^\dagger(\mathbf{k}) H_{mf}(\mathbf{k}) \Psi(\mathbf{k}) + Z \Delta^2 N - 3\lambda N, \quad (7)$$

with $H_{mf}(\mathbf{k}) = \lambda \Omega_0 + 2Z \Delta \gamma_{\mathbf{k}} \Omega_2$. The corresponding Lagrangian is thus given by

$$L_{mf} = \frac{1}{2} \sum_{\mathbf{k}} \Psi^\dagger(\mathbf{k}, i\omega_n) [i\omega_n \Omega_1 - H_{mf}(\mathbf{k})] \Psi(\mathbf{k}, i\omega_n).$$

Here the generalized 8×8 Dirac matrices have been defined by $\Omega_1 = \sigma_z \otimes \sigma_0 \otimes \sigma_0$, $\Omega_2 = \sigma_x \otimes \sigma_z \otimes \sigma_z$, where σ_x , σ_y , σ_z , and σ_0 are three Pauli 2×2 matrices and unity matrix, respectively. Ω_0 is the 8×8 unity matrix, Ω_1 and Ω_2 obey the anticommutation relation $[\Omega_1, \Omega_2]_+ = 2\Omega_0$. From the Lagrangian, the bosonic Matsubara Green function is given by

$$G(\mathbf{k}, i\omega_n) = \frac{-i\omega_n \Omega_1 - \lambda \Omega_0 + 2Z \Delta \gamma_{\mathbf{k}} \Omega_2}{\omega_n^2 + [\lambda^2 - (2Z \Delta \gamma_{\mathbf{k}})^2]}, \quad (8)$$

which implies that the bosonic quasiparticle excitations form a continuum band, and their dispersion relation is $\omega_{\mathbf{k}} = \sqrt{\lambda^2 - (2Z \Delta \gamma_{\mathbf{k}})^2}$. Moreover, the free energy can be evaluated as

$$F_{mf} = \frac{1}{\beta} \sum_{\mathbf{k}, \mu} \ln \left[2 \sinh \left(\frac{\beta \omega_{\mathbf{k}}}{2} \right) \right] + Z \Delta^2 N - 3\lambda N, \quad (9)$$

and the saddle point equations are derived as follows

$$\begin{aligned} \frac{1}{N} \sum_{\mathbf{k}} \frac{2\lambda}{\sqrt{\lambda^2 - (2Z \Delta \gamma_{\mathbf{k}})^2}} [2n_B(\omega_{\mathbf{k}}) + 1] &= 3, \\ \frac{1}{N} \sum_{\mathbf{k}} \frac{(2Z \Delta \gamma_{\mathbf{k}})^2}{\sqrt{\lambda^2 - (2Z \Delta \gamma_{\mathbf{k}})^2}} [2n_B(\omega_{\mathbf{k}}) + 1] &= Z. \end{aligned} \quad (10)$$

The first equation corresponds to the request of the local constraint on average $\sum_{\mu} \langle b_{i, \mu}^\dagger b_{i, \mu} \rangle = 1$, the second one is to determine the short-ranged VB order parameter self-consistently.

To get the collective excitations of the model, the dynamical correlation functions of the spin, orbital, and combined spin-orbital density operators should be calculated. The corresponding density operators can be written in terms of the Nambu spinor introduced above,

$$\begin{aligned} S_i^z &= \frac{1}{4} \Psi^\dagger(\mathbf{r}_i) \Omega_3 \Psi(\mathbf{r}_i), \\ T_i^z &= \frac{1}{4} \Psi^\dagger(\mathbf{r}_i) \Omega_4 \Psi(\mathbf{r}_i), \\ L_i^{zz} &= \frac{1}{4} \Psi^\dagger(\mathbf{r}_i) \Omega_5 \Psi(\mathbf{r}_i), \end{aligned}$$

where $\Omega_3 = \sigma_z \otimes \sigma_0 \otimes \sigma_z$, $\Omega_4 = \sigma_z \otimes \sigma_z \otimes \sigma_0$, and $\Omega_5 = \sigma_0 \otimes \sigma_z \otimes \sigma_z$ are introduced. Except for the anticommutation relations $[\Omega_1, \Omega_2]_+ = [\Omega_2, \Omega_3]_+ = [\Omega_2, \Omega_4]_+ = 2\Omega_0$, all other relations between the Dirac matrices from Ω_1 to Ω_5 satisfy the commutation relations.

In 1D system, we have $Z = 2$ and $\gamma_k = \cos k$. At $T = 0\text{K}$, there are no particle excitations $n_B(\omega_k) = 0$. When the summations over momenta are converted into the integrals within the first Brillouin zone, the self-consistent equations are easily solved, leading to $\Delta \simeq 1.67476$ and $\lambda \simeq 7.31391$. The corresponding ground state energy per site is given by $\varepsilon_g = -2\Delta^2 \simeq -5.60964$, and there is a finite gap in the bosonic quasiparticle excitation spectrum, $\omega_0 = \sqrt{\lambda^2 - 16\Delta^2} \simeq 2.93533$ at the momentum $k = 0$ and $k = \pi$. These results have shown that the ground state of the model is a quantum *disordered* state of resonating valence bonds, in qualitative agreement with the previous numerical calculations [16]. Moreover, the dynamical susceptibilities at zero temperature can be easily evaluated as follows

$$\begin{aligned} \chi(q, \omega + i\eta) &= \int \frac{dk}{16} \left[\frac{\lambda^2 \pm 16\Delta^2 \cos k \cos(k+q)}{\omega_k \omega_{k+q}} - 1 \right] \\ &\times \left[\frac{1}{\omega + \omega_k + \omega_{k+q} + i\eta} - \frac{1}{\omega - \omega_k - \omega_{k+q} + i\eta} \right], \end{aligned} \quad (11)$$

where η is a positive infinitesimal, both the spin and orbital dynamic susceptibilities are equal to each other and choose the lower “-” sign, and the combined spin-orbital dynamic susceptibility takes the upper “+” sign. Then, the dynamic density spectra are easily obtained

$$\begin{aligned} \text{Im } \chi_S(q \rightarrow Q, \omega) &= \text{Im } \chi_T(q \rightarrow Q, \omega) = \text{Im } \chi_L(q \rightarrow 0, \omega) \\ &= \frac{1}{8|\omega|} \sqrt{\frac{4\lambda^2 - \omega^2}{\omega^2 - 4\lambda^2 + 64\Delta^2}} \end{aligned} \quad (12)$$

where $2\sqrt{\lambda^2 - 16\Delta^2} < |\omega| < 2\lambda$. Clearly, there is a finite gap $E_r = 2\omega_0 \simeq 5.87066$ in both the spin and orbital collective excitations at $q = Q$, and in the combined spin-orbital collective excitation at $q = 0$. Therefore we conclude that the ground state is a quantum disordered spin-orbital liquid state of short-ranged valence bonds with a finite gap in their collective excitations.

On a 2D square lattice, $Z = 4$ and $\gamma_{\mathbf{k}} = \frac{1}{2}(\cos k_x + \cos k_y)$. At zero temperature, the conversion from the summations over momenta to the integrals will be *invalid* as $(8\Delta) \rightarrow \lambda$. Following the analogous treatments for the Bose-Einstein condensation [22], we separate the divergent terms at $\mathbf{k}^* = \mathbf{0}$ and $\mathbf{k}^* = \mathbf{Q}$ from the summations to yield

$$\int \frac{d^2k}{(2\pi)^2} \frac{\lambda}{\sqrt{\lambda^2 - (8\Delta\gamma_{\mathbf{k}})^2}} + \frac{4n_B(\omega_{\mathbf{k}^*})}{\sqrt{1 - (8\Delta/\lambda)^2}} = \frac{3}{2},$$

$$\int \frac{d^2k}{(2\pi)^2} \frac{\lambda\gamma_{\mathbf{k}}^2}{\sqrt{\lambda^2 - (8\Delta\gamma_{\mathbf{k}})^2}} + \frac{4n_B(\omega_{\mathbf{k}^*})}{\sqrt{1 - (8\Delta/\lambda)^2}} = \frac{\lambda}{16}. \quad (13)$$

When $(8\Delta) \rightarrow \lambda$, the boson condensation happens, and the ratio $\rho = 4n_B(\omega_{\mathbf{k}^*})/\sqrt{1 - (8\Delta/\lambda)^2}$ is finite, which can be determined from the first equation by setting $\lambda = 8\Delta$ inside the integral: $\rho \simeq 0.107$. Similarly, from the second equation one can also obtain the short-ranged VB order parameter $\Delta \simeq 1.3159$ and thus $\lambda \simeq 10.5271$. Moreover, the ground state energy per site is also evaluated to be $\varepsilon_g = -4\Delta^2 \simeq -6.9263$. The bosonic quasi-particle excitation spectrum now becomes linearly dependence of momentum near the minimal points $\mathbf{k}^* = \mathbf{0}$ and $\mathbf{k}^* = \mathbf{Q}$. When the boson condensation is carefully treated, the dynamical structure factors for spin, orbital, and combined spin-orbital collective excitations can also be calculated through the fluctuation dissipation theorem. After some algebra, the final results are given by

$$S_S(\mathbf{q} \rightarrow \mathbf{Q}, \omega) = S_T(\mathbf{q} \rightarrow \mathbf{Q}, \omega) = S_L(\mathbf{q} \rightarrow \mathbf{0}, \omega)$$

$$= \frac{\pi\rho^2}{2}\delta(\omega)\text{sign}\omega + \frac{1}{\pi\lambda}K\left(\frac{|\omega|}{2\lambda}\right)\sqrt{\left(\frac{2\lambda}{\omega}\right)^2 - 1}, \quad (14)$$

where $K(x)$ is the complete elliptical function of the second kind. In the range of $0 < |\omega| < 2\lambda$, there is a sharp resonance in the collective excitations at zero frequency, corresponding to the formation of a novel long-range ordered spin-orbital crystal state of resonating valence bonds. In such a state, both the spin and orbital degrees of freedom form an AFM long-range order separately, while the combined spin-orbital degrees of freedom has an FM long-range ordering. The corresponding magnetizations can be read off from the coefficient of the delta function $m = \rho/\sqrt{8\pi} \simeq 0.021$, which is much *smaller* than that of the SU(2) antiferromagnetic Heisenberg spin model. Actually, this result is consistent with the identity in Eq.(3), as the VB state in the mean field theory has kept the SU(4) symmetry of the model. The present results are *in conflict with* the prediction of a spin liquid state with a finite energy gap by the quantum Monte Carlo Green function calculations on a *small* square lattice for the same model [17]. However, the very small magnetization clearly indicates that the quantum fluctuations in the spin-orbital coupled model are much stronger than those in the pure spin model. The numerical data shows that the energy gap (even if it exists) is very close to zero. In this sense, both analytical and numerical results seem to be consistent with each other.

Several remarks are in order. i) The reduced model Hamiltonian has displayed a unique form of attractive pairing instability among the nearest neighbor hard-core bosons. Introducing a short-ranged VB pairing order parameter does not break the global SU(4) symmetry of the model. ii) The mean field theory in 1D has given rise to

the correct ground state of a quantum disordered spin-orbital liquid state, which was consistent with numerical results [16]. iii) On a 2D square lattice, the long-range spin-orbital ferromagnetic ordered state is obtained due to the Bose-Einstein condensation. We can also expect that such a long-ranged ordered state should be robust to the perturbations added even away from the SU(4) symmetric point slightly. iv) Compared with the numerical method, the SU(4) Schwinger boson mean field approach for the symmetrically coupled spin-orbital systems has provided the insight on the exact nature of the ground state and its first excitations.

In conclusion, we derive a reduced effective spin-orbital Hamiltonian in a symmetric spin-orbital systems by introducing the SU(4) Schwinger boson representation. After introducing a symmetric pairing VB order parameter, a mean field theory has been developed, leading to a quantum disordered liquid state with a finite gap in the collective excitations in 1D, and on a 2D square lattice a novel long-range FM ordered spin-orbital crystal state where both the spin and orbital degrees of freedom form AFM. The new ordering properties due to the interplay between the spin and orbital degrees of freedom can be detected in the future experiments on transitional metal oxides.

This work was supported by a RGC grant of Hong Kong. G. -M. Zhang thanks Prof. Lu Yu for stimulating his interest in this subject, and acknowledges the partial support from NSF-China (Grant No. 10074036) and the Special Fund for Major State Basic Research Projects of China (G2000067107).

-
- [1] W. Bao, *et al.*, Phys. Rev. Lett. **78**, 507 (1997); C. Broholm, *et al.*, J. Appl. Phys. **79**, 5023 (1996).
 - [2] T. Mizokawa, *et al.*, Phys. Rev. B **61**, R3776 (2000).
 - [3] G. Khaliullin and S. Maekawa, Phys. Rev. Lett. **85**, 3950 (2000).
 - [4] E. Saitoh, *et al.*, Nature **410**, 180 (2001).
 - [5] K. I. Kugel and D. I. Khomskii, Usp. Fiz. Nauk **136**, 621 (1982).
 - [6] L. F. Feiner, *et al.*, Phys. Rev. Lett. **78**, 2799 (1997).
 - [7] D. P. Arovas and A. Auerbach, Phys. Rev. B **52**, 10114 (1995).
 - [8] S. K. Pati and R. R. P. Singh, Phys. Rev. Lett. **81**, 5406 (1998).
 - [9] J. van den Brink, *et al.*, Phys. Rev. B **58**, 10276 (1998).
 - [10] Y. Q. Li, *et al.*, Phys. Rev. Lett. **81**, 3527 (1998).
 - [11] B. Frischmuth, *et al.*, Phys. Rev. Lett. **82**, 835 (1998).
 - [12] A. Joshi, *et al.*, Phys. Rev. B **60**, 6584 (1999).
 - [13] P. Azaria, *et al.*, Phys. Rev. Lett. **83**, 624 (1999).
 - [14] Y. Yamashita, *et al.*, J. Phys. Soc. Japan **69**, 242 (2000).
 - [15] C. Itoi, *et al.*, Phys. Rev. B **61**, 6747 (2000).
 - [16] G. Santoro, *et al.*, Phys. Rev. Lett. **74**, 4039 (1995); Phys. Rev. B **53**, 828 (1996); *ibid.* **55**, 16168 (1997).

- [17] G. Santoro, *et al.*, Phys. Rev. Lett. **83**, 3065 (1999).
- [18] P. Paul, *et al.*, J. Am. Chem. Soc. **116**, 4145 (1994).
- [19] J. Choi, *et al.*, Phys. Rev. Lett. **80**, 1328 (1998); J. Choi, *et al.*, Phys. Rev. B **59**, 1819 (1999).
- [20] D. P. Arovas and A. Auerbach, Phys. Rev. B **38**, 316 (1988); A. Auerbach and D. P. Arovas, Phys. Rev. Lett. **61**, 617 (1988).
- [21] J. E. Hirsch and S. Tang, Phys. Rev. B **39**, 2850 (1989).
- [22] See, for example, K. Huang, *Statistical Mechanics* (McGraw-Hill, New York, 1956).